

# ALGEBRAIC MANIPULATION

## Highest Common Factor (H.C.F.)

If two or more algebraic expressions are given then their common factor of highest power is called the H.C.F of the expressions.

## Least Common Multiple (L.C.M)

If an algebraic expression  $p(x)$  is exactly divisible by two or more expressions, then  $p(x)$  is called the Common Multiple of the given expressions. The Least Common Multiple (L.C.M) is the product of common factors together with non-common factors of the given expressions.

## Finding H.C.F

We can find H.C.F of given expressions by the following two methods.

(i) By Factorization

(ii) By division

## H.C.F. by Factorization

### Example

Find the H.C.F of the following polynomials.

$$x^2 - 4, x^2 + 4x + 4, 2x^2 + x - 6$$

### Solution

By factorization,

$$x^2 - 4 = (x + 2)(x - 2)$$

$$x^2 + 4x + 4 = (x + 2)^2 = (x + 2)(x + 2)$$

$$2x^2 + x - 6 = 2x^2 + 4x - 3x - 6 = 2x(x + 2) - 3(x + 2) \\ = (x + 2)(2x - 3)$$

Common factors =  $x + 2$

$$\text{H.C.F} = x + 2$$

## H.C.F. by Division

### Example

Use division method to find the H.C.F. of the polynomials

$$p(x) = x^3 - 7x^2 + 14x - 8 \text{ and}$$

$$q(x) = x^3 - 7x + 6$$

### Solution

$$\begin{array}{r} x^3 - 7x + 6 \overline{) x^3 - 7x^2 + 14x - 8} \\ \underline{+ x^3 \quad \quad - 7x + 6} \phantom{- 8} \\ -7x^2 + 21x - 14 \end{array}$$

Here the remainder can be factorized as

$$-7x^2 + 21x - 14 = -7(x^2 - 3x + 2)$$

We ignore  $-7$  because it is not common to both the given polynomials and consider  $x^2 - 3x + 2$ .

$$\begin{array}{r} x^2 - 3x + 2 \overline{) x^3 + 0x^2 - 7x + 6} \\ \underline{+ x^3 - 3x^2 + 2x} \phantom{+ 6} \\ 3x^2 - 9x + 6 \\ \underline{3x^2 - 9x + 6} \\ 0 \end{array}$$

Hence H.C. F of  $p(x)$  and  $q(x)$  is  $x^2 - 3x + 2$

**Example**

Find the L.C.M of  $p(x)=12(x^3-y^3)$  and  $q(x)=8(x^3-xy^2)$

**Solution**

By prime factorization of the given expressions, we have

$$p(x)=12(x^3-y^3)=2^2 \times 3 \times (x-y)(x^2+xy+y^2) \text{ and}$$

$$q(x)=8(x^3-xy^2)=8x(x^2-y^2)=2^3 x(x+y)(x-y) \text{ Hence L.C.M. of } p(x) \text{ and } q(x),$$

$$2^3 \times 3 \times x(x+y)(x-y)(x^2+xy+y^2)=24x(x+y)(x^3-y^3)$$

**Relation between H.C.F and L.C.M****Example**

By factorization, find (i) H.C.F (ii) L.C.M of  $p(x)=12(x^5-x^4)$  and  $q(x)=8(x^4-3x^3+3x^2)$ . Establish a relation between  $p(x)$ ,  $q(x)$  and H.C.F and L.C.M of the expressions  $p(x)$  and  $q(x)$ .

**Solution**

Firstly, let us factorize completely the given expressions  $p(x)$  and  $q(x)$  into irreducible factors. We have

$$p(x)=12(x^5-x^4)=12x^4(x-1)=2^2 \times 3 \times x^4(x-1) \text{ and}$$

$$q(x) = 8(x^4-3x^3+2x^2)=8x^2(x^2-3x+2)=2^3 x^2(x-1)(x-2)$$

$$\text{H.C.F. of } p(x) \text{ and } q(x) = 2^2 x^2(x-1)=4x^2(x-1)$$

$$\text{L.C.M of } p(x) \text{ and } q(x) = 2^3 \times 3 \times x^4(x-1)(x-2)$$

$$\begin{aligned} \text{Now } p(x) \times q(x) &= 12x^4(x-1) \times 8x^2(x-1)(x-2) \\ &= 96x^6(x-1)^2(x-2) \dots\dots\dots(i) \end{aligned}$$

and (L.C.M) (H.C.F)

$$= [2^3 \times 3 \times x^4(x-1)(x-2)] [4x^2(x-1)]$$

$$= [24x^4(x-1)(x-2)] [4x^2(x-1)]$$

$$= 96x^6(x-1)^2(x-2) \dots\dots\dots(ii)$$

From (i) and (ii)

$$\text{L.C.M} \times \text{H.C.F} = P(x) \times q(x)$$

**Note**

$$(1) \quad \text{L.C.M} = \frac{p(x) \times q(x)}{\text{H.C.F}} \quad \text{or}$$

$$\text{H.C.F} = \frac{p(x) \times q(x)}{\text{L.C.M}}$$

(2) If L.C.M, H.C.F and one of  $p(x)$  or  $q(x)$  are known, then

$$p(x) = \frac{\text{L.C.M} \times \text{H.C.F}}{q(x)}$$

$$q(x) = \frac{\text{L.C.M} \times \text{H.C.F}}{p(x)}$$

**Example**

Find H.C.F of the polynomials,

$$p(x) = 20(2x^3 + 3x^2 - 2x)$$

$$q(x) = 9(5x^4 + 40x)$$

Then using the above formula (I) find the L.C.M of  $p(x)$  and  $q(x)$ .

**Solution**

We have

$$p(x) = 20(2x^3 + 3x^2 - 2x) = 20x(2x^2 + 3x - 2)$$

$$= 20x(2x^2 + 4x - x - 2) = 20x[2x(x+2) - (x+2)] = 20x(x+2)(2x-1) = 2^2 \times 5 \times x(x+2)(2x-1)$$

$$q(x) = 9(5x^4 + 40x) = 45x(x^3 + 8) = 45x\left[(x^3) + (2)^3\right]$$

$$= 45x(x+2)(x^2 - 2x + 4) = 5 \times 3^2 \times x(x+2)(x^2 - 2x + 4) \text{ Thus H.C.F of } p(x) \text{ and } q(x) \text{ is:}$$

$$= 5x(x+2)$$

$$\text{Now, using the formula} \quad \text{L.C.M.} = \frac{p(x) \times q(x)}{\text{H.C.F}}$$

We obtain

$$\begin{aligned} \text{L.C.M.} &= \frac{2^2 \times 5 \times x(x+2)(2x-1) \times 5 \times 3^2 \times x(x+2)(x^2 - 2x + 4)}{5x(x+2)} \\ &= 4 \times 5 \times 9 \times x(x+2)(2x-1)(x^2 - 2x + 4) \\ &= 180x(x+2)(2x-1)(x^2 - 2x + 4) \end{aligned}$$

**Example**

Find the L.C.M of

$$p(x) = 6x^3 - 7x^2 - 27x + 8 \quad \text{and}$$

$$q(x) = 6x^3 + 17x^2 + 9x - 4$$

**Solution**

We have, by long division,

$$\begin{array}{r} 1 \\ 6x^3 - 7x^2 - 27x + 8 \overline{) 6x^3 + 17x^2 + 9x - 4} \\ \underline{6x^3 - 7x^2 - 27x + 8} \phantom{0} \\ 24x^2 + 36x - 12 \end{array}$$

But the remainder  $24x^2 + 36x - 12$

$$= 12(2x^2 + 3x - 1)$$

Thus, ignoring 12, we have

$$\begin{array}{r} 2x^2 + 3x - 1 \overline{) 6x^3 - 7x^2 - 27x + 8} \\ \underline{6x^3 + 9x^2 - 3x} \phantom{0} \\ -16x^2 - 24x + 8 \\ \underline{-16x^2 - 24x + 8} \phantom{0} \\ 0 \end{array}$$

Hence H.C.F of  $p(x)$  and  $q(x)$  is

$$= 2x^2 + 3x - 1$$

$$x^2 + 6x - 27 = x^2 - 3x + 9x - 27$$

$$= x(x-3) + 9(x-3)$$

$$= (x-3)(x+9) \quad \dots\dots(ii)$$

$$2x^2 - 18 = 2(x^2 - 9)$$

$$= 2[(x)^2 - (3)^2]$$

$$= 2(x+3)(x-3) \quad \dots\dots(iii)$$

From (i), (ii) and (iii)

Common factors =  $(x-3)$

$$HCF = x-3$$

$$iii) x^3 - 2x^2 + x, x^2 + 2x - 3, x^2 + 3x - 4$$

**Sol:** By factorization

$$x^3 - 2x^2 + x = x(x^2 - 2x + 1)$$

$$= x(x^2 - x - x + 1)$$

$$= x[x(x-1) - 1(x-1)]$$

$$= x(x-1)(x-1) \quad \dots\dots(i)$$

$$x^2 + 2x - 3 = x^2 - x + 3x - 3$$

$$= x(x-1) + 3(x-1)$$

$$= (x-1)(x+3) \quad \dots\dots(ii)$$

$$x^2 + 3x - 4 = x^2 - x + 4x - 4$$

$$= x(x-1) + 4(x-1)$$

$$= (x-1)(x+4) \quad \dots\dots(iii)$$

From (i), (ii) and (iii)

Common factors:  $x-1$

$$HCF = x-1$$

$$iv) 18(x^3 + 9x^2 + 8x), 24(x^2 - 3x + 2)$$

**Sol:** By factorization

$$18(x^3 + 9x^2 + 8x) = 18x(x^2 + 9x + 8)$$

$$= 18x(x^2 - x - 8x + 8)$$

$$= 18x[x(x-1) - 8(x-1)]$$

$$= 2 \times 3 \times 3 \times x(x-1)(x-8) \quad \dots\dots(i)$$

$$24(x^2 - 3x + 2) =$$

$$24(x^2 - x - 2x + 2)$$

$$= 2 \times 2 \times 2 \times 3[x(x-1) - 2(x-1)]$$

$$= 2 \times 2 \times 2 \times 3(x-1)(x-2) \quad \dots(ii)$$

From (i) and (ii)

$$HCF = 2 \times 3(x-1)$$

$$= 6(x-1)$$

$$v) 36(3x^4 + 5x^3 - 2x^2), 54(27x^4 - x)$$

**Sol:** By factorization

$$36(3x^4 + 5x^3 - 2x^2) = 36x^2(3x^2 + 5x - 2)$$

$$= 36x^2(3x^2 + 6x - x - 2)$$

$$= 36x^2[3x(x+2) - 1(x+2)]$$

$$= 2 \times 2 \times 3 \times 3 \times x \times x(x+2)(3x-1) \quad \dots(i)$$

$$54(27x^4 - x) = 54x(27x^3 - 1)$$

$$= 54x[(3x)^3 - (1)^3]$$

$$= 54x(3x-1)[(3x)^2 + (3x)(1) + (1)^2]$$

$$= 2 \times 3 \times 3 \times 3 \times x(3x-1)(9x^2 + 3x + 1) \quad \dots(ii)$$

From (i) and (ii)

$$\text{Common factors} = 2, 3, 3, x, (3x-1)$$

$$HCF = 2 \times 3 \times 3 \times x(3x-1)$$

$$= 18x(3x-1)$$

**Q3. Find the H.C.F of the following by division methal.**

$$i) p(x) = x^3 + 3x^2 - 16x + 12, q(x) = x^3 + x^2 - 10x + 8$$

$$\begin{array}{r} \text{Sol: } x^3 + x^2 - 10x + 8 \overline{) x^3 + 3x^2 - 16x + 12} \\ \underline{-x^3 + x^2 + 10x + 8} \\ 2x^2 - 6x + 4 \end{array}$$

Dividing remainder by 2



$$\begin{array}{r}
 x^2 - 3x + 2 \\
 x^2 - 3x + 2 \overline{) x^6 + x^2 - 10x + 8} \\
 \underline{-x^6 + 3x^2 + 2x} \phantom{+ 8} \\
 4x^2 - 12x + 8 \\
 \underline{-4x^2 + 12x - 8} \\
 0
 \end{array}$$

Hence HCF =  $x^2 - 3x + 2$

ii)  $P(x) = x^4 + x^3 - 2x^2 + x - 3$ ,  
 $q(x) = 5x^3 + 3x^2 - 17x + 6$

$$\begin{array}{r}
 5x^3 + 3x^2 - 17x + 6 \overline{) x^4 + x^3 - 2x^2 + x - 3} \\
 \underline{5x^4 + 5x^3 - 10x^2 + 5x - 15} \quad \text{(Multiplying by 5)} \\
 -5x^4 + 3x^3 + 17x^2 + 6x \phantom{- 3} \\
 \underline{2x^3 + 7x^2 - x - 15} \\
 \underline{10x^3 + 35x^2 - 5x - 75} \quad \text{(Multiplying by 5)} \\
 -10x^3 + 6x^2 + 34x + 12 \\
 \hline
 29x^2 + 29x - 87
 \end{array}$$

Divided by 29

$$x^2 + x - 3$$

$$\begin{array}{r}
 5x - 2 \\
 x^2 + x - 3 \overline{) 5x^3 + 3x^2 - 17x + 6} \\
 \underline{-5x^3 + 5x^2 + 15x} \phantom{+ 6} \\
 -2x^2 - 2x + 6 \\
 \underline{-2x^2 + 2x + 6} \\
 0
 \end{array}$$

Hence H.C.F =  $x^2 + x - 3$

iii)  $p(x) = 2x^5 - 4x^4 - 6x$ ,  
 $q(x) = x^5 + x^4 - 3x^3 - 3x^2$

$$\begin{array}{r}
 2 \\
 x^5 + x^4 - 3x^3 - 3x^2 \overline{) 2x^5 - 4x^4 - 6x} \\
 \underline{-2x^5 + 2x^4} \phantom{+ 6x} \\
 -6x^4 + 6x^3 + 6x^2 - 6x
 \end{array}$$

Dividing by -6

$$\begin{array}{r}
 x^4 - x^3 - x^2 + x \\
 x^4 - x^3 - x^2 + x \overline{) x^6 + x^4 - 3x^3 - 3x^2} \\
 \underline{-x^6 + x^4 + x^3 + x^2} \phantom{+ 2x} \\
 2x^4 - 2x^3 - 4x^2 \\
 \underline{-2x^4 + 2x^3 + 2x^2 + 2x} \\
 -2x^2 - 2x
 \end{array}$$

Dividing by -2

$$x^2 + x$$

$$\begin{array}{r}
 x^2 - 2x + 1 \\
 x^2 + x \overline{) x^4 - x^3 - x^2 + x} \\
 \underline{-x^4 + x^3} \phantom{+ x} \\
 -2x^3 - x^2 + x \\
 \underline{-2x^3 + 2x^2} \phantom{+ x} \\
 x^2 + x \\
 \underline{-x^2 - x} \\
 0
 \end{array}$$

Hence H.C.F =  $x^2 + x = x(x+1)$

**Q4. Find the L.C.M of the following expressions:**

i)  $39x^7y^3z$  and  $91x^5y^6z^7$

**Sol:** By factorization

$$39x^7y^3z = 13 \times 3 \times x \times x \times x \times x \times x \times y \times y \times y \times z$$

$$91x^5y^6z^7 = 13 \times 7 \times x \times x \times x \times x \times y \times y \times y \times y \times y \times z \times z \times z \times z \times z \times z$$

Hence L.C.M =

$$13 \times 3 \times 7 \times x \times x \times x \times x \times x \times y \times y \times y \times y \times y \times z \times z \times z \times z \times z \times z = 273x^7y^6z^7$$

ii)  $102xy^2z$ ,  $85x^2yz$  and  $187xyz^2$

**Sol:** By factorization

$$102xy^2z = 2 \times 3 \times 17 \times x \times y \times y \times z$$

$$85x^2yz = 5 \times 17 \times x \times x \times y \times z$$

$$187xyz^2 = 11 \times 17 \times x \times y \times z \times z$$

$$\begin{aligned}\text{Hence L.C.M} &= 17 \times 11 \times 5 \times 3 \times 2 \times x \times x \times y \times y \times z \times z \\ &= 5610x^2y^2z^2\end{aligned}$$

**Q5. Find the L.C.M of the following expressions by factorization:**

i)  $x^2 - 25x + 100$  and  $x^2 - x - 20$

**Sol:** By factorization

$$\begin{aligned}x^2 - 25x + 100 &= x^2 - 5x - 20x + 100 \\ &= x(x-5) - 20(x-5) \\ &= (x-5)(x-20) \dots\dots\dots(i) \\ x^2 - x - 20 &= x^2 - 5x + 4x - 20 \\ &= x(x-5) + 4(x-5) \\ &= (x-5)(x+4) \dots\dots\dots(ii)\end{aligned}$$

From (i) and (ii)

$$\text{L.C.M} = (x-5)(x-20)(x+4)$$

ii)  $x^2 + 4x + 4$ ,  $x^2 - 4$ ,  $2x^2 + x - 6$

**Sol:** By factorization

$$\begin{aligned}x^2 + 4x + 4 &= x^2 + 2x + 2x + 4 \\ &= x(x+2) + 2(x+2) \\ &= (x+2)(x+2) \dots\dots\dots(i) \\ x^2 - 4 &= (x)^2 - (2)^2 \\ &= (x+2)(x-2) \dots\dots\dots(ii) \\ 2x^2 + x - 6 &= 2x^2 + 4x - 3x - 6 \\ &= 2x(x+2) - 3(x+2) \\ &= (x+2)(2x-3) \dots\dots\dots(iii)\end{aligned}$$

From (i), (ii) and (iii)

$$\begin{aligned}\text{LCM} &= (x+2)(x+2)(x-2)(2x-3) \\ &= (x+2)^2(x-2)(2x-3)\end{aligned}$$

iii)  $2(x^4 - y^4)$ ,  $3(x^3 + 2x^2y - xy^2 - 2y^3)$

**Sol:** By factorization

$$2(x^4 - y^4) = 2[(x^2)^2 - (y^2)^2]$$

$$\begin{aligned}&= 2(x^2 + y^2)(x^2 - y^2) \\ &= 2(x^2 + y^2)(x+y)(x-y) \dots\dots\dots(i)\end{aligned}$$

$$\begin{aligned}3(x^3 + 2x^2y - xy^2 - 2y^3) &= 3[x^2(x+2y) - y^2(x+2y)] \\ &= 3(x+2y)(x^2 - y^2) \\ &= 3(x+2y)(x+y)(x-y) \dots\dots\dots(ii)\end{aligned}$$

From (i) & (ii)

L.C.M =

$$\begin{aligned}&2 \times 3(x+y)(x-y)(x^2 + y^2)(x+2y) \\ &= 6(x^4 - y^4)(x+2y)\end{aligned}$$

iv)  $4(x^4 - 1)$ ,  $6(x^3 - x^2 - x + 1)$

**Sol:** By factorization

$$\begin{aligned}4(x^4 - 1) &= 4[(x^2)^2 - (1)^2] \\ &= 4(x^2 + 1)(x^2 - 1) \\ &= 2 \times 2(x^2 + 1)[(x)^2 - (1)^2] \\ &= 2 \times 2(x^2 + 1)(x+1)(x-1) \dots\dots\dots(i) \\ 6(x^3 - x^2 - x + 1) &= 6[x^2(x-1) - 1(x-1)] \\ &= 6(x-1)(x^2 - 1) = 2 \times 3(x-1)[(x)^2 - (1)^2] \\ &= 2 \times 3(x-1)(x-1)(x+1) \dots\dots(ii)\end{aligned}$$

From (i) & (ii)

$$\begin{aligned}\text{LCM} &= 2 \times 2 \times 3(x+1)(x-1)(x^2 + 1)(x-1) \\ &= 12(x^4 - 1)(x-1)\end{aligned}$$

**Q6. For what value of  $k$  is  $(x+4)$ , the H.C.F of  $x^2 + x - (2k+2)$  and  $2x^2 + kx - 12$ ?**

**Sol:**  $k = ?$

$$p(x) = x^2 + x - (2k+2) \text{ and}$$

$$q(x) = 2x^2 + kx - 12$$

As given that  $x+4$  is HCF, so  $p(x)$  and  $q(x)$  will be exactly divisible by  $(x+4)$

$$\begin{array}{r}
 x-3 \\
 x+4 \overline{) x^2 + x - (2k+2)} \\
 \underline{\cancel{x^2} + 4x} \phantom{- (2k+2)} \\
 -3x - (2k+2) \\
 \underline{\phantom{-} 3x + 12} \\
 12 - (2k+2)
 \end{array}$$

$$= 12 - 2k - 2$$

$$= 10 - 2k$$

As  $p(x)$  is exactly divisible by  $x+4$ , so,

$$10 - 2k = 0$$

$$10 = 2k$$

$$\frac{10}{2} = k$$

$$k = 5$$

**Q7.** If  $(x+3)(x-2)$  is the H.C.F of

$p(x) = (x+3)(2x^2 - 3x + k)$  and

$q(x) = (x-2)(3x^2 + 7x - l)$ , find  $k$  and  $l$ .

**Sol:**  $k = ?$  and  $l = ?$

As  $(x+3)(x-2)$  is the H.C.F, so  $p(x)$  and  $q(x)$  will be exactly divisible by

$(x+3)(x-2)$  i.e.,  $\frac{p(x)}{HCF}$  has remainder zero.

$$\begin{array}{r}
 \frac{(x+3)(2x^2 - 3x + k)}{(x+3)(x-2)} = \frac{2x^2 - 3x + k}{x-2} \\
 \text{i.e.} \quad \begin{array}{r}
 2x+1 \\
 x-2 \overline{) 2x^2 - 3x + k} \\
 \underline{\pm 2x^2 \mp 4x} \phantom{+ k} \\
 x + k \\
 \underline{\phantom{x} \pm x \mp 2} \\
 k + 2
 \end{array}
 \end{array}$$

As remainder = 0, then

$$k + 2 = 0$$

$$\boxed{k = -2}$$

and  $\frac{q(x)}{HCF}$  has zero remainder

$$\frac{(x-2)(3x^2 + 7x - l)}{(x+3)(x-2)} = \frac{3x^2 + 7x - l}{x+3}$$

$$\begin{array}{r}
 3x-2 \\
 x+3 \overline{) 3x^2 + 7x - l} \\
 \underline{\pm 3x^2 \pm 9x} \phantom{- l} \\
 -2x - l \\
 \underline{\phantom{-} 2x \mp 6} \\
 -l + 6
 \end{array}$$

As remainder = 0

$$-l + 6 = 0$$

$$-l = -6$$

$$\Rightarrow \boxed{l = 6}$$

**Q8.** The LCM and HCF of two polynomials  $p(x)$  and  $q(x)$  are  $2(x^4 - 1)$  and  $(x+1)(x^2 + 1)$  respectively. If  $p(x) = x^3 + x + 1$ , find  $q(x)$ .

**Sol:** LCM =  $2(x^4 - 1)$ ,

$$HCF = (x+1)(x^2 + 1)$$

$$p(x) = x^3 + x^2 + x + 1, \quad q(x) = ?$$

$$\text{As } p(x) \times q(x) = (LCM) \times (HCF)$$

$$q(x) = \frac{(LCM) \times (HCF)}{p(x)}$$

$$= \frac{2(x^4 - 1) \times (x+1)(x^2 + 1)}{x^3 + x^2 + x + 1}$$

$$= \frac{2(x^4 - 1)(x^3 + x^2 + x + 1)}{x^3 + x^2 + x + 1}$$

$$q(x) = 2(x^4 - 1)$$



**Q9.** Let  $p(x) = 10(x^2 - 9)(x^2 - 3x + 2)$  and  $q(x) = 10x(x+3)(x-1)^2$ . If the H.C.F. of  $p(x), q(x)$  is  $10(x+3)(x-1)$ , find their L.C.M.

**Sol:**  $p(x) = 10(x^2 - 9)(x^2 - 3x + 2)$ ,

$$q(x) = 10x(x+3)(x-1)^2$$

$$\text{H.C.F.} = 10(x+3)(x-1), \text{ L.C.M.} = ?$$

$$\text{As } (L.C.M.) \times (H.C.F.) = p(x) \times q(x)$$

$$\text{L.C.M.} = \frac{p(x) \times q(x)}{\text{H.C.F.}}$$

$$= \frac{10(x^2 - 9)(x^2 - 3x + 2) \times 10x(x+3)(x-1)^2}{10(x+3)(x-1)}$$

$$= \frac{(x^2 - 9)(x^2 - 3x + 2) \times 10x \cancel{(x+3)} \cancel{(x-1)} (x-1)}{\cancel{(x+3)} \cancel{(x-1)}}$$

$$= 10x(x-1)(x^2 - 9)(x^2 - 3x + 2)$$

$$= 10x(x-1)(x^2 - 9)(x^2 - x - 2x + 2)$$

$$= 10x(x-1)(x^2 - 9)[x(x-1) - 2(x-1)]$$

$$= 10x(x-1)(x^2 - 9)(x-1)(x-2)$$

$$= 10x(x-1)^2(x^2 - 9)(x-2)$$

**Q10.** Let the product of L.C.M and H.C.F of two polynomials be  $(x+3)^2(x-2)(x+5)$ . If one polynomial is  $(x+3)(x-2)$  and the second polynomial is  $x^2 + kx + 15$ , find the value of  $k$ .

**Sol:**  $k = ?$

Product of L.C.M. & H.C.F is

$$\text{LCM} \times \text{HCF} = (x+3)^2(x-2)(x+5)$$

$$p(x) = (x+3)(x-2)$$

$$q(x) = x^2 + kx + 15$$

$$\text{As } p(x) \times q(x) = \text{LCM} \times \text{HCF}$$

$$(x+3)(x-2)(x^2 + kx + 15)$$

$$= (x+3)^2(x-2)(x+5)$$

$$x^2 + kx + 15 = \frac{(x+3)\cancel{(x+3)}\cancel{(x-2)}(x+5)}{\cancel{(x+3)}\cancel{(x-2)}}$$

$$x^2 + kx + 15 = (x+3)(x+5)$$

$$x^2 + kx + 15 = x^2 + 3x + 5x + 15$$

$$x^2 + kx + 15 = x^2 + 8x + 15$$

Comparing co-efficient of 'x'

$$\Rightarrow kx = 8x$$

$$\boxed{k = 8}$$

**Q11.** Waqas wishes to distribute 128 bananas and also 176 apples equally among a certain number of children. Find the highest number of the Children. Who can get the fruit in this way?

**Sol:** No. of bananas = 128

No. of apples = 176

Highest no. of children who get the fruit in this way is H.C.F.

So No. of bananas =

$$2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2$$

No. of apples =

$$2 \times 2 \times 2 \times 2 \times 11$$

Hence required no. of children =

$$2 \times 2 \times 2 \times 2 = 16$$

### Example

Simplify

$$\frac{x+3}{x^2-3x+2} + \frac{x+2}{x^2-4x+3} + \frac{x+1}{x^2-5x+6}, x \neq 1, 2, 3$$

### Solution

$$\frac{x+3}{x^2-3x+2} + \frac{x+2}{x^2-4x+3} + \frac{x+1}{x^2-5x+6}$$

$$= \frac{x+3}{x^2-2x-x+2} + \frac{x+2}{x^2-3x-x+3} + \frac{x+1}{x^2-3x-2x+6}$$

$$= \frac{x+3}{x(x-2)-1(x-2)} + \frac{x+2}{x(x-3)-1(x-3)} + \frac{x+1}{x(x-3)-2(x-3)}$$



$$\begin{aligned}
 &= \frac{x+3}{(x-2)(x-1)} + \frac{x+2}{(x-3)(x-1)} + \frac{x+1}{(x-3)(x-2)} \\
 &= \frac{(x+3)(x-3) + (x+2)(x-2) + (x+1)(x-1)}{(x-1)(x-2)(x-3)} \\
 &= \frac{x^2-9+x^2-4+x^2-1}{(x-1)(x-2)(x-3)} \\
 &= \frac{3x^2-14}{(x-1)(x-2)(x-3)}
 \end{aligned}$$

### Example

Express the product  $\frac{x^3-8}{x^2-4} \times \frac{x^2+6x+8}{x^2-2x+1}$

as an algebraic expression reduced lowest forms  $x \neq 2, -2, 1$

### Solution

By factorizing completely, we have

$$\begin{aligned}
 &\frac{x^3-8}{x^2-4} \times \frac{x^2+6x+8}{x^2-2x+1} \\
 &= \frac{(x-2)(x^2+2x+4) \times (x+2)(x+4)}{(x-2)(x+2) \times (x-1)^2} \dots (i)
 \end{aligned}$$

Now the factors of numerator are  $(x-2), (x^2+2x+4), (x+2)$  and  $(x+4)$  and the factors of denominator are  $(x-2), (x+2)$  and  $(x-1)^2$ .

Therefore, their H.C.F. is  $(x-2) \times (x+2)$

By cancelling H.C.F i.e.,  $(x-2) \times (x+2)$  from (i), we get the simplified form of given product as the fraction  $\frac{(x^2+2x+4)(x+4)}{(x-1)^2}$

### Example

Divide  $\frac{x^2+x+1}{x^2-9}$  by  $\frac{x^3-1}{x^2-4x+3}$

and simplify by reducing to lowest forms.

### Solution

$$\begin{aligned}
 &\text{We have } \frac{x^2+x+1}{x^2-9} \div \frac{x^3-1}{x^2-4x+3} \\
 &= \frac{(x^2+x+1)}{(x^2-9)} \times \frac{(x^2-4x+3)}{(x^3-1)} \\
 &= \frac{(x^2+x+1)(x^2-x-3x+3)}{(x^2-9)(x^3-1)} \\
 &= \frac{(x^2+x+1)[x(x-1)-3(x-1)]}{(x+3)(x-3)(x-1)(x^2+x+1)} \\
 &= \frac{(x^2+x+1)(x-3)(x-1)}{(x+3)(x-3)(x-1)(x^2+x+1)} = \frac{1}{x+3}, x \neq -3
 \end{aligned}$$

## Exercise 6.2

Simplify each of the following as a rational expression.

Q1. 
$$\begin{aligned}
 &\frac{x^2-x-6}{x^2-9} + \frac{x^2+2x-24}{x^2-x-12} \\
 &= \frac{x^2-3x+2x-6}{(x)^2-(3)^2} + \frac{x^2+6x-4x-24}{x^2+3x-4x-12}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{x(x-3)+2(x-3)}{(x+3)(x-3)} + \frac{x(x+6)-4(x+6)}{x(x+3)-4(x+3)} \\
 &= \frac{(x-3)(x+2)}{(x+3)(x-3)} + \frac{(x+6)(x-4)}{(x+3)(x-4)} \\
 &= \frac{x+2}{x+3} + \frac{x+6}{x+3} = \frac{x+2+x+6}{x+3}
 \end{aligned}$$

$$= \frac{2x+8}{x+3}$$

$$= \frac{2(x+4)}{x+3}$$

Q2.  $\left[ \frac{x+1}{x-1} - \frac{x-1}{x+1} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1}$

$$= \left[ \frac{(x+1)^2 - (x-1)^2}{(x-1)(x+1)} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1}$$

$$= \left[ \frac{(x^2+2x+1) - (x^2-2x+1)}{(x)^2 - (1)^2} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1}$$

$$= \left[ \frac{x^2+2x+1-x^2+2x-1}{x^2-1} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1}$$

$$= \left[ \frac{4x}{x^2-1} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1}$$

$$= \left[ \frac{4x(x^2+1) - 4x(x^2-1)}{(x^2-1)(x^2+1)} \right] + \frac{4x}{x^4-1}$$

$$= \frac{4x^3+4x-4x^3+4x}{(x^2)^2 - (1)^2} + \frac{4x}{x^4-1}$$

$$= \frac{8x}{x^4-1} + \frac{4x}{x^4-1}$$

$$= \frac{8x+4x}{x^4-1}$$

$$= \frac{12x}{x^4-1}$$

Q3.  $\frac{1}{x^2-8x+15} + \frac{1}{x^2-4x+3} - \frac{2}{x^2-6x+5}$

$$= \frac{1}{x^2-3x-5x+15} + \frac{1}{x^2-3x-x+3} - \frac{2}{x^2-5x-x+5}$$

$$= \frac{1}{x(x-3)-5(x-3)} + \frac{1}{x(x-3)-1(x-3)} - \frac{2}{x(x-5)-1(x-5)}$$

$$= \frac{1}{(x-3)(x-5)} + \frac{1}{(x-3)(x-1)} - \frac{2}{(x-5)(x-1)}$$

$$= \frac{x-1+x-5-2(x-3)}{(x-1)(x-3)(x-5)}$$

$$= \frac{x-1+x-5-2x+6}{(x-1)(x-3)(x-5)}$$

$$= \frac{2x-6-2x+6}{(x-1)(x-3)(x-5)}$$

$$= \frac{0}{(x-1)(x-3)(x-5)}$$

$$= 0$$

Q4.  $\frac{(x+2)(x+3)}{x^2-9} + \frac{(x+2)(2x^2-32)}{(x-4)(x^2-x-6)}$

$$= \frac{(x+2)(x+3)}{(x)^2 - (3)^2} + \frac{(x+2) \cdot 2(x^2-16)}{(x-4)(x^2+2x-3x-6)}$$

$$= \frac{(x+2)(x+3)}{(x-3)(x+3)} + \frac{2(x+2)[(x)^2 - (4)^2]}{(x-4)(x^2+2x-3x-6)}$$

$$= \frac{(x+2)}{x-3} + \frac{2(x+2)(x+4)(x-4)}{(x-4)(x+2)(x-3)}$$

$$= \frac{x+2}{x-3} + \frac{2x+8}{x-3}$$

$$= \frac{x+2+2x+8}{x-3}$$

$$= \frac{3x+10}{x-3}$$

Q5.  $\frac{x+3}{2x^2+9x+9} + \frac{1}{2(2x-3)} - \frac{4x}{4x^2-9}$

$$= \frac{x+3}{2x^2+6x+3x+9} + \frac{1}{2(2x-3)} - \frac{4x}{(2x)^2 - (3)^2}$$

$$= \frac{x+3}{2x(x+3)+3(x+3)} + \frac{1}{2(2x-3)} - \frac{4x}{(2x+3)(2x-3)}$$

$$\begin{aligned}
&= \frac{\cancel{x+3}}{(\cancel{x+3})(2x+3)} + \frac{1}{2(2x-3)} - \frac{4x}{(2x+3)(2x-3)} \\
&= \frac{1}{2x+3} + \frac{1}{2(2x-3)} - \frac{4x}{(2x+3)(2x-3)} \\
&= \frac{2(2x-3) + 2x+3 - 2(4x)}{2(2x+3)(2x-3)} \\
&= \frac{4x-6+2x+3-8x}{2(2x+3)(2x-3)} \\
&= \frac{-2x-3}{2(2x+3)(2x-3)} \\
&= \frac{-1(\cancel{2x+3})}{2(\cancel{2x+3})(2x-3)} \\
&= \frac{-1}{2(2x-3)} \\
&= \frac{1}{2(3-2x)}
\end{aligned}$$

**Q6.**  $A - \frac{1}{A}$ , where  $A = \frac{a+1}{a-1}$

$$\text{so } \frac{1}{A} = \frac{a-1}{a+1}$$

$$\begin{aligned}
\text{Now } A - \frac{1}{A} &= \frac{a+1}{a-1} - \frac{a-1}{a+1} \\
&= \frac{(a+1)^2 - (a-1)^2}{(a-1)(a+1)} \\
&= \frac{(a^2 + 2a + 1) - (a^2 - 2a + 1)}{(a)^2 - (1)^2} \\
&= \frac{\cancel{a^2} + 2a + \cancel{1} - \cancel{a^2} + 2a - \cancel{1}}{a^2 - 1} \\
&= \frac{4a}{a^2 - 1}
\end{aligned}$$

**Q7.**  $\left[ \frac{x-1}{x-2} + \frac{2}{2-x} \right] - \left[ \frac{x+1}{x+2} + \frac{4}{4-x^2} \right]$

$$\begin{aligned}
&= \left[ \frac{-(x-1)}{2-x} + \frac{2}{2-x} \right] - \left[ \frac{x+1}{x+2} + \frac{4}{(2)^2 - (x)^2} \right] \\
&= \left[ -\frac{(x-1)}{2-x} + \frac{2}{2-x} \right] - \left[ \frac{x+1}{x+2} + \frac{4}{(2+x)(2-x)} \right] \\
&= \left[ \frac{-x+1+2}{2-x} \right] - \left[ \frac{(x+1)(2-x)+4}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \left[ \frac{2x-x^2+2-x+4}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \left[ \frac{6+x-x^2}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \left[ \frac{6+3x-2x-x^2}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \left[ \frac{3(2+x)-x(2+x)}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \left[ \frac{(\cancel{2+x})(3-x)}{(\cancel{2+x})(2-x)} \right] \\
&= \frac{3-x}{2-x} - \frac{3-x}{2-x} \\
&= \frac{3-x-3+x}{2-x} \\
&= \frac{0}{2-x} \\
&= 0
\end{aligned}$$

**Q8.** What rational expression should be subtracted from  $\frac{2x^2+2x-7}{x^2+x-6}$  to get

$$\frac{x-1}{x-2} = ?$$



**Sol:** Let the required expression be A,

$$\text{then } \frac{2x^2 + 2x - 7}{x^2 + x - 6} - A = \frac{x-1}{x-2}$$

$$\text{or } \frac{2x^2 + 2x - 7}{x^2 + x - 6} - \frac{x-1}{x-2} = A$$

$$\begin{aligned} \text{So } A &= \frac{2x^2 + 2x - 7}{x^2 + 3x - 2x - 6} - \frac{x-1}{x-2} \\ &= \frac{2x^2 + 2x - 7}{x(x+3) - 2(x+3)} - \frac{x-1}{x-2} \\ &= \frac{2x^2 + 2x - 7}{(x+3)(x-2)} - \frac{x-1}{x-2} \\ &= \frac{2x^2 + 2x - 7 - (x-1)(x+3)}{(x+3)(x-2)} \\ &= \frac{2x^2 + 2x - 7 - (x^2 - x + 3x - 3)}{(x+3)(x-2)} \\ &= \frac{(2x^2 + 2x - 7) - (x^2 + 2x - 3)}{(x+3)(x-2)} \\ &= \frac{2x^2 + 2x - 7 - x^2 - 2x + 3}{(x+3)(x-2)} \\ &= \frac{x^2 - 4}{(x+3)(x-2)} \\ &= \frac{(x)^2 - (2)^2}{(x+3)(x-2)} \\ &= \frac{(x+2)(x-2)}{(x+3)(x-2)} \\ &= \frac{x+2}{x+3} \end{aligned}$$

**Perform the indicated operations and simplify to the lowest forms.**

$$\text{Q9. } \frac{x^2 + x - 6}{x^2 - x - 6} \times \frac{x^2 - 4}{x^2 - 9}$$

$$\begin{aligned} &= \frac{x^2 + 3x - 2x - 6}{x^2 - 3x + 2x - 6} \times \frac{(x)^2 - (2)^2}{(x)^2 - (3)^2} \\ &= \frac{x(x+3) - 2(x+3)}{x(x-3) + 2(x-3)} \times \frac{(x+2)(x-2)}{(x+3)(x-3)} \\ &= \frac{(x+3)(x-2)}{(x-3)(x+2)} \times \frac{(x+2)(x-2)}{(x+3)(x-3)} \\ &= \frac{(x-2)^2}{(x-3)^2} \end{aligned}$$

$$\begin{aligned} \text{Q10. } &\frac{x^3 - 8}{x^2 - 4} \times \frac{x^2 + 6x + 8}{x^2 - 2x + 1} \\ &= \frac{(x)^3 - (2)^3}{(x)^2 - (2)^2} \times \frac{x^2 + 2x + 4x + 8}{x^2 - x - x + 1} \\ &= \frac{(x-2)[(x)^2 + (x)(2) + (2)^2]}{(x-2)(x+2)} \times \frac{x(x+2) + 4(x+2)}{x(x-1) - 1(x-1)} \\ &= \frac{x^2 + 2x + 4}{x+2} \times \frac{(x+2)(x+4)}{(x-1)(x-1)} \\ &= \frac{(x^2 + 2x + 4)(x+4)}{(x-1)^2} \end{aligned}$$

$$\begin{aligned} \text{Q11. } &\frac{x^4 - 8x}{2x^2 + 5x - 3} \times \frac{2x-1}{x^2 + 2x + 4} \times \frac{x+3}{x^2 - 2x} \\ &= \frac{x(x^3 - 8)}{2x^2 + 6x - x - 3} \times \frac{2x-1}{x^2 + 2x + 4} \times \frac{x+3}{x(x-2)} \\ &= \frac{x[(x)^3 - (2)^3]}{2x(x+3) - 1(x+3)} \times \frac{2x-1}{x^2 + 2x + 4} \times \frac{x+3}{x(x-2)} \\ &= \frac{x(x-2)(x^2 + 2x + 4)}{(x+3)(2x-1)} \times \frac{2x-1}{x^2 + 2x + 4} \times \frac{x+3}{x(x-2)} \\ &= 1 \end{aligned}$$

$$\text{Q12. } \frac{2y^2 + 7y - 4}{3y^2 - 13y + 4} \div \frac{4y^2 - 1}{6y^2 + y - 1}$$

$$\begin{aligned}
&= \frac{2y^2 + 8y - y - 4}{3y^2 - y - 12y + 4} \div \frac{(2y)^2 - (1)^2}{6y^2 + 3y - 2y - 1} \\
&= \frac{2y(y+4) - 1(y+4)}{y(3y-1) - 4(3y-1)} \div \frac{(2y+1)(2y-1)}{3y(2y+1) - 1(2y+1)} \\
&= \frac{(y+4)\cancel{(2y-1)}}{\cancel{(3y-1)}(y-4)} \div \frac{\cancel{(2y+1)}(2y-1)}{\cancel{(2y+1)}\cancel{(3y-1)}} \\
&= \frac{(y+4)\cancel{(2y-1)}}{\cancel{(3y-1)}(y-4)} \times \frac{\cancel{(2y+1)}(3y-1)}{\cancel{(2y+1)}\cancel{(2y-1)}} \\
&= \frac{y+4}{y-4}
\end{aligned}$$

**Q13.**  $\left[ \frac{x^2 + y^2}{x^2 - y^2} - \frac{x^2 - y^2}{x^2 + y^2} \right] \div \left[ \frac{x+y}{x-y} - \frac{x-y}{x+y} \right]$

$$\begin{aligned}
&= \left[ \frac{(x^2 + y^2)^2 - (x^2 - y^2)^2}{(x^2 - y^2)(x^2 + y^2)} \right] \div \left[ \frac{(x+y)^2 - (x-y)^2}{(x-y)(x+y)} \right] \\
&= \frac{x^4 + y^4 + 2x^2y^2 - (x^4 + y^4 - 2x^2y^2)^2}{(x^2 - y^2)(x^2 + y^2)} \\
&\quad + \frac{x^2 + y^2 + 2xy - x^2 - y^2 + 2xy}{x^2 - y^2} \\
&= \frac{\cancel{x^4} + \cancel{y^4} + 2x^2y^2 - \cancel{x^4} - \cancel{y^4} + 2x^2y^2}{(x^2 - y^2)(x^2 + y^2)} \\
&\quad + \frac{\cancel{x^2} + \cancel{y^2} + 2xy - \cancel{x^2} - \cancel{y^2} + 2xy}{x^2 - y^2} \\
&= \frac{4x^2y^2}{(x^2 - y^2)(x^2 + y^2)} \div \frac{4xy}{x^2 - y^2} \\
&= \frac{\cancel{4x^2}\cancel{y^2}}{(\cancel{x^2} - \cancel{y^2})(x^2 + y^2)} \times \frac{\cancel{x^2} - \cancel{y^2}}{\cancel{4xy}} \\
&= \frac{xy}{x^2 + y^2}
\end{aligned}$$

### Square Root of Algebraic Expression

The square root of a given expression  $p(x)$  as another expression  $q(x)$  such that  $q(x) \cdot q(x) = p(x)$ .

As  $5 \times 5 = 25$ , so square root of 25 is 5

It means we can find square root of the expression  $p(x)$  if it can be expressed as a perfect square.

#### Example

Use factorization to find the square root of the expression

$$4x^2 - 12x + 9$$

#### Solution

$$\begin{aligned}
&\text{We have, } 4x^2 - 12x + 9 \\
&= 4x^2 - 6x - 6x + 9 = 2x(2x-3) - 3(2x-3) \\
&= (2x-3)(2x-3) = (2x-3)^2
\end{aligned}$$

$$\begin{aligned}
&\text{Hence } \sqrt{4x^2 - 12x + 9} \\
&= \pm(2x-3)
\end{aligned}$$

#### Example

Find the square root of

$$x^2 + \frac{1}{x^2} + 12\left(x + \frac{1}{x}\right) + 38, x \neq 0$$

#### Solution

$$\begin{aligned}
&\text{We have } x^2 + \frac{1}{x^2} + 12\left(x + \frac{1}{x}\right) + 38 \\
&= x^2 + \frac{1}{x^2} + 2 + 12\left(x + \frac{1}{x}\right) + 36, \\
&\quad (\text{adding and subtracting } 2)
\end{aligned}$$

$$= \left[ \pm \left( x + \frac{1}{x} + 6 \right) \right]^2;$$

Hence the required square root is

$$\pm \left( x + \frac{1}{x} + 6 \right)$$

Find the square root of  $4x^4 + 12x^3 + x^2 - 12x + 4$

$$\begin{array}{r}
 2x^2 + 3x - 2 \\
 \hline
 2x^2 \phantom{+ 3x - 2} \\
 \hline
 4x^2 + 3x \phantom{- 2} \\
 \hline
 4x^2 + 6x - 2 \\
 \hline
 -8x^2 - 12x + 4 \\
 \hline
 +8x^2 + 12x + 4 \\
 \hline
 0
 \end{array}$$

Thus square root of given expression is  $\pm(2x^2+3x-2)$

Find the square root of the expression

$$4\frac{x^2}{y^2}+8\frac{x}{y}+16+12\frac{y}{x}+9\frac{y^2}{x^2}$$

We note that the given expression is in descending powers of  $x$ .

$$\begin{array}{r}
 2\frac{x}{y} + 2 + 3\frac{y}{x} \\
 \hline
 4\frac{x^2}{y^2} + 8\frac{x}{y} + 16 + 12\frac{y}{x} + 9\frac{y^2}{x^2} \\
 \hline
 2\frac{x}{y} \\
 \hline
 4\frac{x}{y} + 2 \\
 \hline
 4\frac{x}{y} + 4 + 3\frac{y}{x} \\
 \hline
 12 + 12\frac{y}{x} + 9\frac{y^2}{x^2} \\
 \hline
 \pm 12 \pm 12\frac{y}{x} \pm 9\frac{y^2}{x^2} \\
 \hline
 0
 \end{array}$$

Hence the square root of given expression is  $\pm \left( 2\frac{x}{y} + 2 + 3\frac{y}{x} \right)$

To make the expression  $x^4 - 10x^3 + 33x^2 - 42x + 20$  a perfect square,

- What should be added to it?
- What should be subtracted from it?
- What should be the value of  $x$ ?

$x^2$	$x^2 - 5x + 4$ <hr/> $x^4 - 10x^3 + 33x^2 - 42x + 20$ <hr/> $+ x^4$
$2x^2 - 5x$	$-10x^3 + 33x^2$ <hr/> $-10x^3 + 25x^2$ <hr/> $+$
$2x^2 - 10x + 4$	$8x^2 - 42x + 20$ <hr/> $-8x^2 - 40x + 16$ <hr/> $+$ <hr/> $-2x + 4$ <hr/>

For making the given expression a perfect square the remainder must be zero.



Hence

(i) We should add  $(2x-4)$  to the given expression

(ii) We should subtract  $(-2x+4)$  from the given expression

### Exercise 6.3

**Q1.** Use factorization to find the square root of the following expressions.

$$\begin{aligned} \text{i)} \quad & 4x^2 - 12xy + 9y^2 \\ &= (2x)^2 - 2(2x)(3y) + (3y)^2 \\ &= (2x - 3y)^2 \end{aligned}$$

$$\begin{aligned} \text{Hence } & \sqrt{4x^2 - 12xy + 9y^2} \\ &= \sqrt{(2x - 3y)^2} \\ &= \pm(2x - 3y) \end{aligned}$$

$$\begin{aligned} \text{ii)} \quad & x^2 - 1 + \frac{1}{4x^2} \\ &= (x)^2 - 2(x)\left(\frac{1}{2x}\right) + \left(\frac{1}{2x}\right)^2 \end{aligned}$$

$$\begin{aligned} \text{Hence } & \sqrt{x^2 - 1 + \frac{1}{4x^2}} \\ &= \sqrt{\left(x - \frac{1}{2x}\right)^2} \\ &= \pm\left(x - \frac{1}{2x}\right) \end{aligned}$$

$$\begin{aligned} \text{iii)} \quad & \frac{1}{16}x^2 - \frac{1}{12}xy + \frac{1}{36}y^2 \\ &= \left(\frac{1}{4}x\right)^2 - 2\left(\frac{1}{4}x\right)\left(\frac{1}{6}y\right) + \left(\frac{1}{6}y\right)^2 \end{aligned}$$

(iii) We should take  $-2x+4=0$  to find the value of  $x$ . This gives the required value of  $x$  i.e.,  $x=2$ .

$$= \left(\frac{1}{4}x - \frac{1}{6}y\right)^2$$

$$\begin{aligned} \text{Hence } & \sqrt{\frac{1}{16}x^2 - \frac{1}{12}xy + \frac{1}{36}y^2} \\ &= \sqrt{\left(\frac{1}{4}x - \frac{1}{6}y\right)^2} \\ &= \pm\left(\frac{1}{4}x - \frac{1}{6}y\right) \end{aligned}$$

$$\begin{aligned} \text{iv)} \quad & 4(a+b)^2 - 12(a^2 - b^2) + 9(a-b)^2 \\ &= [2(a+b)]^2 - 2 \times 2(a+b) \times 3(a-b) + [3(a-b)]^2 \\ &= [2(a+b) - 3(a-b)]^2 \\ &= (-a + 5b)^2 \\ &= (5b - a)^2 \end{aligned}$$

$$\begin{aligned} \text{Hence } & \sqrt{4(a+b)^2 - 12(a^2 - b^2) + 9(a-b)^2} \\ &= \sqrt{(5b - a)^2} \\ &= \pm(5b - a) \end{aligned}$$

$$\begin{aligned} \text{v)} \quad & \frac{4x^6 - 12x^3y^3 + 9y^6}{9x^4 + 24x^2y^2 + 16y^4} \\ &= \frac{(2x^3)^2 - 2(2x^3)(3y^3) + (3y^3)^2}{(3x^2)^2 + 2(3x^2)(4y^2) + (4y^2)^2} \end{aligned}$$

$$= \frac{(2x^3 - 3y^3)^2}{(3x^2 + 4y^2)^2}$$

$$\begin{aligned} \text{Hence } \sqrt{\frac{4x^6 - 12x^3y^3 + 9y^6}{9x^4 + 24x^2y^2 + 16y^4}} \\ = \sqrt{\frac{(2x^3 - 3y^3)^2}{(3x^2 + 4y^2)^2}} \\ = \pm \left( \frac{2x^3 - 3y^3}{3x^2 + 4y^2} \right) \end{aligned}$$

$$\begin{aligned} \text{vi) } \left(x + \frac{1}{x}\right)^2 - 4\left(x - \frac{1}{x}\right) \quad (x \neq 0) \\ = (x)^2 + \left(\frac{1}{x}\right)^2 + 2\left(\cancel{x}\right)\left(\cancel{\frac{1}{x}}\right) - 4\left(x - \frac{1}{x}\right) \\ = x^2 + \frac{1}{x^2} + 2 - 4\left(x - \frac{1}{x}\right) \dots\dots\dots(i) \end{aligned}$$

$$\text{Let } x - \frac{1}{x} = a$$

$$\text{Squaring } \left(x - \frac{1}{x}\right)^2 = (a)^2$$

$$x^2 + \frac{1}{x^2} - 2 = a^2$$

$$x^2 + \frac{1}{x^2} = a^2 + 2$$

So expression (i) becomes

$$= a^2 + 2 + 2 - 4a$$

$$= a^2 - 4a + 4$$

$$= (a)^2 - 2(a)(2) + (2)^2$$

$$= (a - 2)^2$$

Putting value of 'a'

$$= \left(x - \frac{1}{x} - 2\right)^2$$

$$\text{Hence } = \sqrt{\left(x - \frac{1}{x} - 2\right)^2}$$

$$= \pm \left(x - \frac{1}{x} - 2\right)$$

$$\text{vii) } \left(x^2 + \frac{1}{x^2}\right)^2 - 4\left(x + \frac{1}{x}\right)^2 + 12 \dots(i)$$

$$\text{Let } x + \frac{1}{x} = a$$

$$\text{Squaring } \left(x + \frac{1}{x}\right)^2 = (a)^2$$

$$x^2 + \frac{1}{x^2} + 2 = a^2$$

$$x^2 + \frac{1}{x^2} = a^2 - 2$$

So expression (i) becomes

$$= (a^2 - 2)^2 - 4(a)^2 + 12$$

$$= (a^2)^2 - 2(a^2)(2) + (2)^2 - 4a^2 + 12$$

$$= a^4 - 4a^2 + 4 - 4a^2 + 12$$

$$= a^4 - 8a^2 + 16$$

$$= (a^2)^2 - 2(a^2)(4) + (4)^2$$

$$= (a^2 - 4)^2$$

Putting values of  $a^2$

$$= \left(x^2 + \frac{1}{x^2} + 2 - 4\right)^2$$

$$= \left(x^2 + \frac{1}{x^2} - 2\right)^2$$

$$\text{Hence } = \sqrt{\left(x^2 + \frac{1}{x^2}\right)^2 - 4\left(x + \frac{1}{x}\right)^2 + 12}$$

$$= \sqrt{\left(x^2 + \frac{1}{x^2} - 2\right)^2}$$

$$= \pm \left( x^2 + \frac{1}{x^2} - 2 \right)$$

$$\begin{aligned} \text{viii)} \quad & (x^2 + 3x + 2)(x^2 + 4x + 3)(x^2 + 5x + 6) \\ &= (x^2 + x + 2x + 2)(x^2 + x + 3x + 3)(x^2 + 2x + 3x + 6) \\ &= [x(x+1) + 2(x+1)][x(x+1) + 3(x+1)][x(x+2) + 3(x+2)] \\ &= (x+1)(x+2)(x+1)(x+3)(x+2)(x+3) \\ &= (x+1)^2(x+2)^2(x+3)^2 \end{aligned}$$

Hence

$$\begin{aligned} & \sqrt{(x^2 + 3x + 2)(x^2 + 4x + 3)(x^2 + 5x + 6)} \\ &= \sqrt{(x+1)^2(x+2)^2(x+3)^2} \\ &= \pm (x+1)(x+2)(x+3) \end{aligned}$$

$$\begin{aligned} \text{ix)} \quad & (x^2 + 8x + 7)(2x^2 - x - 3)(2x^2 + 11x - 21) \\ &= (x^2 + x + 7x + 7)(2x^2 + 2x - 3x - 3)(2x^2 + 14x - 3x - 21) \\ &= [x(x+1) + 7(x+1)][2x(x+1) - 3(x+1)] \\ & \quad [2x(x+7) - 3(x+7)] \\ &= (x+1)(x+7)(x+1)(2x-3)(x+7)(2x-3) \\ &= (x+1)^2(x+7)^2(2x-3)^2 \end{aligned}$$

Hence

$$\begin{aligned} & \sqrt{(x^2 + 8x + 7)(2x^2 - x - 3)(2x^2 + 11x - 21)} \\ &= \sqrt{(x+1)^2(x+7)^2(2x-3)^2} \\ &= \pm (x+1)(x+7)(2x-3) \end{aligned}$$

**Q2.** Use division method to find the square root of the following expressions.

$$\text{i)} \quad 4x^2 + 12xy + 9y^2 + 16x + 24y + 16$$

	$2x + 3y + 4$
$2x$	$4x^2 + 12xy + 9y^2 + 16x + 24y + 16$ $\underline{4x^2}$
$4x + 3y$	$12xy + 9y^2 + 16x + 24y + 16$ $\underline{12xy + 9y^2}$
$4x + 6y + 4$	$16x + 24y + 16$ $\underline{16x + 24y + 16}$
	0

Hence the square root of given expression is  
 $\pm (2x + 3y + 4)$

$$\text{ii)} \quad x^4 - 10x^3 + 37x^2 - 60x + 36$$

	$x^2 - 5x + 6$
$x^2$	$x^4 - 10x^3 + 37x^2 - 60x + 36$ $\underline{-x^4}$
$2x^2 - 5x$	$-10x^3 + 37x^2 - 60x + 36$ $\underline{+10x^3 \pm 25x^2}$
$2x^2 - 10x + 6$	$-12x^2 - 60x + 36$ $\underline{-12x^2 \mp 60x \pm 36}$
	0

Hence  $\sqrt{x^4 - 10x^3 + 37x^2 - 60x + 36}$   
 $= \pm (x^2 - 5x + 6)$

$$\text{iii)} \quad 9x^4 - 6x^3 + 7x^2 - 2x + 1$$



$$\begin{array}{r}
 3x^2 - x + 1 \\
 3x^2 \overline{) 9x^4 - 6x^3 + 7x^2 - 2x + 1} \\
 \underline{-9x^4} \phantom{+ 7x^2 - 2x + 1} \\
 6x^3 - x^2 + 7x^2 - 2x + 1 \\
 \underline{-6x^3 + x^2} \phantom{- 2x + 1} \\
 6x^2 - 2x + 1 \\
 \underline{-6x^2 + 2x} \phantom{+ 1} \\
 0
 \end{array}$$

Hence  $\sqrt{9x^4 - 6x^3 + 7x^2 - 2x + 1}$   
 $= \pm(3x^2 - x + 1)$

iv)  $4 + 25x^2 - 12x - 24x^3 + 16x^4$   
 In descending order  
 $= 16x^4 - 24x^3 + 25x^2 - 12x + 4$

$$\begin{array}{r}
 4x^2 - 3x + 2 \\
 4x^2 \overline{) 16x^4 - 24x^3 + 25x^2 - 12x + 4} \\
 \underline{-16x^4} \phantom{+ 25x^2 - 12x + 4} \\
 8x^3 - 3x^2 + 25x^2 - 12x + 4 \\
 \underline{-8x^3 + 9x^2} \phantom{- 12x + 4} \\
 8x^2 - 6x + 2 \\
 \underline{-8x^2 + 12x} \phantom{+ 2} \\
 0
 \end{array}$$

Hence  $\sqrt{16x^4 - 24x^3 + 25x^2 - 12x + 4}$   
 $= \pm(4x^2 - 3x + 2)$

v)  $\frac{x^2}{y^2} - 10\frac{x}{y} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2}$   
 $(x \neq 0, y \neq 0)$

Hence

$$\begin{array}{r}
 \frac{x}{y} - 5 + \frac{y}{x} \\
 \frac{x}{y} \overline{) \frac{x^2}{y^2} - 10\frac{x}{y} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2}} \\
 \underline{-\frac{x^2}{y^2}} \phantom{- 10\frac{x}{y} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2}} \\
 2x - 5y - 10\frac{y}{x} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2} \\
 \underline{-2x + 10\frac{y}{x} - 25} \phantom{+ 27 - 10\frac{y}{x} + \frac{y^2}{x^2}} \\
 2x - 10 + \frac{y}{x} - 10\frac{y}{x} + \frac{y^2}{x^2} \\
 \underline{-2x + 10 - \frac{y}{x} + \frac{y^2}{x^2}} \\
 0
 \end{array}$$

$$\sqrt{\frac{x^2}{y^2} - 10\frac{x}{y} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2}}$$

The required square root

$$= \pm \left( \frac{x}{y} - 5 + \frac{y}{x} \right)$$

Q3. Find the value of 'k' for which the following expression will become a perfect square?

i)  $4x^4 - 12x^3 + 37x^2 - 42x + k$

$$\begin{array}{r}
 2x^2 - 3x + 7 \\
 2x^2 \overline{) 4x^4 - 12x^3 + 37x^2 - 42x + k} \\
 \underline{-4x^4} \phantom{+ 37x^2 - 42x + k} \\
 4x^3 - 3x^2 + 37x^2 - 42x + k \\
 \underline{-4x^3 + 9x^2} \phantom{- 42x + k} \\
 4x^2 - 6x + 7 \\
 \underline{-4x^2 + 12x - 49} \phantom{+ k} \\
 k - 49
 \end{array}$$

As given that the given expression is a perfect square, so

$$\text{Remainder} = 0$$

$$k - 49 = 0$$

$$k = 49$$

ii)  $x^4 - 4x^3 + 10x^2 - kx + 9$

$$\begin{array}{r}
 x^2 \overline{) x^4 - 4x^3 + 10x^2 - kx + 9} \\
 \underline{-x^4} \phantom{+ 10x^2 - kx + 9} \\
 2x^2 - 2x \phantom{+ 9} \\
 \underline{2x^2 - 4x + 3} \\
 (-k+12)x
 \end{array}$$

As given that the given expression is a perfect square, so

$$\text{Remainder} = 0$$

$$(-k+12)x = 0$$

As  $x \neq 0$ , so  $-k+12=0$

$$\Rightarrow k = 12$$

**Q4.** Find the values of 'l' and 'm' for which the following expression will become perfect square.

i)  $x^4 + 4x^3 + 16x^2 + lx + m$

$$\begin{array}{r}
 x^2 \overline{) x^4 + 4x^3 + 16x^2 + lx + m} \\
 \underline{-x^4} \phantom{+ 16x^2 + lx + m} \\
 2x^2 + 2x \phantom{+ m} \\
 \underline{2x^2 + 4x + 6} \\
 (l-24)x + (m-36)
 \end{array}$$

As the given expression is to be a perfect square, so

$$\text{Remainder} = 0$$

$$(l-24)x + (m-36) = 0$$

As  $x \neq 0$ , so  $l-24=0$  and  $m-36=0$

$$\Rightarrow l = 24 \text{ and } m = 36$$

ii)  $49x^4 - 70x^3 + 109x^2 + lx - m$

$$\begin{array}{r}
 7x^2 \overline{) 49x^4 - 70x^3 + 109x^2 + lx - m} \\
 \underline{-49x^4} \phantom{+ 109x^2 + lx - m} \\
 14x^2 - 5x \phantom{- m} \\
 \underline{14x^2 - 10x + 6} \\
 (l+60)x - m - 36
 \end{array}$$

As the given expression is to be a perfect square, so

$$(l+60)x - m - 36 = 0$$

As  $x \neq 0$ , so  $l+60=0$  and  $-m-36=0$

$$\Rightarrow l = -60 \text{ and } m = -36$$

**Q5.** To make the expression

$9x^4 - 12x^3 + 22x^2 - 13x + 12$  a perfect square.

i) What should be added to it?

ii) What should be subtracted from it?

iii) What should be the value of 'x'?

$$\begin{array}{r}
 3x^2 \overline{) 9x^4 - 12x^3 + 22x^2 - 13x + 12} \\
 \underline{-9x^4} \phantom{+ 22x^2 - 13x + 12} \\
 6x^2 - 2x \phantom{+ 12} \\
 \underline{6x^2 - 4x + 3} \\
 -x + 3
 \end{array}$$

To make the given expression a complete square

i)  $x-3$  should be added

ii)  $-x+3$  should be subtracted

iii) For value of 'x'

$$\text{Remainder} = 0$$

$$-x + 3 = 0$$

$$\boxed{x = 3}$$

**Q6. Find H.C.F of following by factorization**

$$8x^4 - 128, 12x^3 - 96.$$

**Solution:**

$$\begin{aligned} 8x^4 - 128 &= 8(x^4 - 16) \\ &= 8((x^2)^2 - (4)^2) \\ &= 8(x^2 + 4)(x^2 - 4) \\ &= 8(x^2 + 4)(x + 2)(x - 2) \\ 12x^3 - 96 &= 12(x^3 - 8) \\ &= 12(x^3 - 2^3) \\ &= 12(x - 2)(x^2 + 2x + 4) \end{aligned}$$

$$\text{Common factor} = 4(x - 2)$$

$$\text{H.C.F} = 4(x - 2)$$

**Q7. Find H.C.F of following by division method.**

$$y^3 + 3y^2 - 3y - 9, y^3 + 3y^2 - 8y - 24$$

**Solution:**

$$\begin{array}{r} 1 \\ y^3 + 3y^2 - 3y - 9 \quad y^3 + 3y^2 - 8y - 24 \\ \hline -y^3 + 3y^2 + 3y + 9 \\ -5y - 15 \\ -5(y + 3) \\ \hline y^2 - 3 \\ (y + 3) \quad y^3 + 3y^2 - 3y - 9 \\ \hline -y^3 + 3y^2 \\ -3y - 9 \\ +3y + 9 \\ \hline x \end{array}$$

$$\text{H.C.F} = y + 3$$

**Q8. Find L.C.M of following by factorization.**

$$12x^2 - 75, 6x^2 - 13x - 5, 4x^2 - 20x + 25$$

**Solution:**

$$\begin{aligned} 12x^2 - 75 &= 3(4x^2 - 25) \\ &= 3((2x)^2 - (5)^2) \\ &= 3(2x + 5)(2x - 5) \\ 6x^2 - 13x - 5 &= 6x^2 - 15x + 2x - 5 \\ &= 3x(2x - 5) + 1(2x - 5) \end{aligned}$$

$$\begin{aligned} &= (3x + 1)(2x - 5) \\ 4x^2 - 20x + 25 &= (2x)^2 + (5)^2 - 2(2x)(5) \\ &= (2x - 5)^2 \\ &= (2x - 5)(2x - 5) \end{aligned}$$

$$\begin{aligned} \text{L.C.M} &= (2x - 5)^2 \times 3(2x + 5)(3x + 1) \\ &= 3(2x - 5)^2(2x + 5)(3x + 1) \end{aligned}$$

**Q9. If H.C.F of  $x^4 + 3x^3 + 5x^2 + 26x + 56$  and  $x^4 + 2x^3 - 4x^2 - x + 28$  is  $x^2 + 5x + 7$ , find the**

**Solution:**

$$\text{L.C.M} = \frac{(x^4 + 3x^3 + 5x^2 + 26x + 56)(x^4 + 2x^3 - 4x^2 - x + 28)}{x^2 + 5x + 7}$$

$$\begin{array}{r} x^2 + 5x + 7 \overline{) x^4 + 3x^3 + 5x^2 + 26x + 56} \\ \underline{-x^4 + 5x^3 + 7x^2} \phantom{+ 26x + 56} \\ -2x^3 - 2x^2 + 26x + 56 \\ \underline{+2x^3 + 10x^2 + 14x} \phantom{+ 56} \\ 8x^2 + 40x + 56 \\ \underline{-8x^2 + 40x + 56} \\ \hline \phantom{0}0 \end{array}$$

**L.C.M**

$$= (x^2 - 2x + 8)(x^4 + 2x^3 - 4x^2 - x + 28)$$

**Q10. Simplify**

$$\begin{aligned} \text{(i)} \quad & \frac{3}{x^3 + x^2 + x + 1} - \frac{3}{x^3 - x^2 + x - 1} \\ &= \frac{3}{(x^2 + 1)(x + 1)} - \frac{3}{(x^2 + 1)(x - 1)} \\ &= \frac{3(x - 1) - 3(x + 1)}{(x^2 + 1)(x + 1)(x - 1)} \\ &= \frac{3x - 3 - 3x - 3}{(x^2 + 1)(x + 1)(x - 1)} \\ &= \frac{-6}{(x^2 + 1)(x + 1)(x - 1)} \\ &= \frac{-6}{(x^2 + 1)(x^2 - 1)} \end{aligned}$$



$$= \frac{-6}{x^4-1} = \frac{6}{1-x^4} \text{ Ans.}$$

$$\begin{aligned} \text{(ii)} \quad & \frac{a+b}{a^2-b^2} \div \frac{a^2-ab}{a^2-2ab+b^2} \\ &= \frac{a+b}{(a-b)(a+b)} \div \frac{a(a-b)}{(a-b)^2} \\ &= \frac{1}{a-b} \div \frac{a}{a-b} \\ &= \frac{1}{\cancel{a-b}} \times \frac{\cancel{a-b}}{a} \\ &= \frac{1}{a} \end{aligned}$$

**Q11. Find square root by using factorization**

$$\left(x^2 + \frac{1}{x^2}\right) + 10\left(x + \frac{1}{x}\right) + 27 \quad (x \neq 0)$$

**Solution:**

$$= \left(x^2 + \frac{1}{x^2}\right) + 10\left(x + \frac{1}{x}\right) + 25 + 2$$

**Q12. Find square root by using division method.**

$$\frac{4x^2}{y^2} + \frac{20x}{y} + 13 - \frac{30y}{x} + \frac{9y^2}{x^2} \quad (x, y \neq 0)$$

**Solution:**

$$\begin{array}{r} \frac{2x}{y} + 5 - \frac{3y}{x} \\ \hline \frac{2x}{y} \quad \frac{4x^2}{y^2} + \frac{20x}{y} + 13 - \frac{30y}{x} + \frac{9y^2}{x^2} \\ \underline{\frac{4x^2}{y^2}} \\ \frac{4x}{y} + 5 \quad \frac{20}{y}x + 13 \\ \underline{-\frac{20}{y}x + 25} \\ \frac{4x}{y} + 10 - \frac{3y}{x} \quad -12 - \frac{30y}{x} + \frac{9y^2}{x^2} \\ \underline{+12 + \frac{30y}{x} + \frac{9y^2}{x^2}} \\ \times \end{array}$$

$$\text{Required square root} = \pm \left( \frac{2x}{y} + 5 - \frac{3y}{x} \right)$$

$$= x^2 + \frac{1}{x^2} + 2 + 10\left(x + \frac{1}{x}\right) + 25$$

$$= \left(x + \frac{1}{x}\right)^2 + 10\left(x + \frac{1}{x}\right) + 25$$

$$\text{Let } x + \frac{1}{x} = a$$

$$= a^2 + 10a + 25$$

$$= (a+5)^2$$

Taking square root

$$= \sqrt{[\pm(a+5)]^2}$$

$$= \pm(a+5)$$

$$= \pm\left(x + \frac{1}{x} + 5\right)$$

## Objective

1. H.C.F of  $p^3q - pq^3$  and  $p^5q^2 - p^2q^5$  is \_\_\_\_  
 (a)  $pq(p^2 - q^2)$  (b)  $pq(p - q)$   
 (c)  $p^2q^2(p - q)$  (d)  $pq(p^3 - q^3)$
2. H.C.F. of  $5x^2y^2$  and  $20x^3y^3$  is: \_\_\_\_  
 (a)  $5x^2y^2$  (b)  $20x^3y^3$   
 (c)  $100x^5y^5$  (d)  $5xy$
3. H.C.F of  $x - 2$  and  $x^2 + x - 6$  is \_\_\_\_  
 (a)  $x^2 + x - 6$  (b)  $x + 2$   
 (c)  $x - 2$  (d)  $x + 2$
4. H.C.F of  $a^3 + b^3$  and  $a^2 - ab + b^2$  is \_\_\_\_  
 (a)  $a + b$   
 (b)  $a^2 - ab + b^2$   
 (c)  $(a - b)^2$  (d)  $a^2 + b^2$
5. H.C.F of  $x^2 - 5x + 6$  and  $x^2 - x - 6$  is \_\_\_\_:  
 (a)  $x - 3$  (b)  $x + 2$   
 (c)  $x^2 - 4$  (d)  $x - 2$
6. H.C.F of  $a^2 - b^2$  and  $a^3 - b^3$  is \_\_\_\_  
 (a)  $a - b$  (b)  $a + b$   
 (c)  $a^2 + ab + b^2$  (d)  $a^2 - ab + b^2$
7. H.C.F of  $x^2 + 3x + 2$ ,  $x^2 + 4x + 3$ ,  $x^2 + 5x + 4$  is:  
 (a)  $x + 1$  (b)  $(x + 1)(x + 2)$   
 (c)  $(x + 3)$  (d)  $(x + 4)(x + 1)$
8. L.C.M of  $15x^2$ ,  $45xy$  and  $30xyz$  is \_\_\_\_  
 (a)  $90xyz$  (b)  $90x^2yz$   
 (c)  $15xyz$  (d)  $15x^2yz$
9. L.C.M of  $a^2 + b^2$  and  $a^4 - b^4$  is: \_\_\_\_  
 (a)  $a^2 + b^2$  (b)  $a^2 - b^2$   
 (c)  $a^4 - b^4$  (d)  $a - b$
10. The product of two algebraic expression is equal to the \_\_\_\_ of

their H.C.F and L.C.M.

- (a) Sum  
 (b) Difference  
 (c) Product  
 (d) Quotient
11. Simplify  $\frac{a}{9a^2 - b^2} + \frac{1}{3a - b} = \text{---}$   
 (a)  $\frac{4a}{9a^2 - b^2}$   
 (b)  $\frac{4a - b}{9a^2 - b^2}$   
 (c)  $\frac{4a + b}{9a^2 - b^2}$   
 (d)  $\frac{b}{9a^2 - b^2}$
12. Simplify  $\frac{a^2 + 5a - 14}{a^2 - 3a - 18} \times \frac{a + 3}{a - 2} = \text{---}$   
 (a)  $\frac{a + 7}{a - 6}$  (b)  $\frac{a + 7}{a - 2}$   
 (c)  $\frac{a + 3}{a - 6}$  (d)  $\frac{a - 3}{a + 2}$
13. Simplify  $\frac{a^3 - b^3}{a^4 - b^4} \div \left( \frac{a^2 + ab + b^2}{a^2 + b^2} \right) = \text{---}$   
 (a)  $\frac{1}{a + b}$  (b)  $\frac{1}{a - b}$   
 (c)  $\frac{a - b}{a^2 + b^2}$  (d)  $\frac{a + b}{a^2 + b^2}$
14. Simplify :  
 $\left( \frac{2x + y}{x + y} - 1 \right) \div \left( 1 - \frac{x}{x + y} \right)$   
 = \_\_\_\_

- (a)  $\frac{x}{x+y}$  (b)  $\frac{x}{x-y}$   
 (c)  $\frac{y}{x}$  (d)  $\frac{x}{y}$
15. The square root of  $a^2 - 2a + 1$  is \_\_\_\_  
 (a)  $\pm(a+1)$  (b)  $\pm(a-1)$   
 (c)  $a-1$  (d)  $a+1$
16. What should be added to complete the square of  $x^4 + 64$ ?  
 (a)  $8x^2$  (b)  $-8x^2$   
 (c)  $16x^2$  (d)  $4x^2$
17. The square root of  $x^4 + \frac{1}{x^4} + 2$  is \_\_\_\_  
 (a)  $\pm\left(x + \frac{1}{x}\right)$  (b)  $\pm\left(x^2 + \frac{1}{x^2}\right)$   
 (c)  $\pm\left(x - \frac{1}{x}\right)$  (d)  $\pm\left(x^2 - \frac{1}{x^2}\right)$
18. The square root of  $4x^2 - 12x + 9$  is:  
 (a)  $\pm(2x - 3)$   
 (b)  $\pm(2x + 3)$   
 (c)  $(2x + 3)^2$   
 (d)  $(2x - 3)^2$

19. L.C.M = \_\_\_\_  
 (a)  $\frac{p(x) \times q(x)}{\text{H.C.F}}$  (b)  $\frac{p(x).q(x)}{\text{L.C.M}}$   
 (c)  $\frac{p(x)}{q(x) \times \text{H.C.F}}$  (d)  $\frac{q(x)}{p(x) \times \text{H.C.F}}$
20. H.C.F. = \_\_\_\_  
 (a)  $\frac{p(x) \times q(x)}{\text{L.C.M}}$  (b)  $\frac{p(x) \times q(x)}{\text{H.C.F}}$   
 (c)  $\frac{p(x)}{q(x) \times \text{L.C.M}}$  (d)  $\frac{\text{L.C.M}}{p(x) \times q(x)}$
21. L.C.M x HCF =  
 (a)  $p(x) \times q(x)$  (b)  $p(x) \times \text{H.C.F}$   
 (c)  $q(x) \times \text{L.C.M}$  (d) None
22. Any unknown expression may be found if \_\_\_\_ of them are known by using the relation  
 $\text{L.C.M} \times \text{H.C.F} = p(x) \times q(x)$   
 (a) Two  
 (b) Three  
 (c) Four  
 (d) None

### ANSWER KEY

1.	a	2.	a	3.	c	4.	b	5.	a
6.	a	7.	a	8.	b	9.	c	10.	c
11.	c	12.	a	13.	a	14.	d	15.	b
16.	c	17.	b	18.	a	19.	a	20.	a
21.	a	22.	b						